Black 4/4/59. (Born - at least 3/2) housed only 26 d.
Once born 5/12 4hrs post ~~to~~ observation. Immediate
~~Three trampled~~ Cares, mechanical wiping. Nipples up.
5/25. Nice mac fat/pink 6/14, 4, 6/15 plus no
removal after wiping. Note eyes "Bug past want settle
down" House 8/3, No ocular lesions. Fat eggs 9/4
popped to 9% 9/29. On 6/12, during electrostorm, 14 more
trampled (first time ever seen ~~3~~ nonetheless this
flock, either). No other damage any place during
Nest storm. On 10/25/59, when I was with them,
There was a Thunderstorm & with a loud thunder
clap about 10 seconds away - about 2 miles away - they
were staring up in the air, the only chickens in the
place to react at Nest time, so.
11/16 - 3 big non-laying pullets to lab. Negative upon
exam for obesity, which was attributed to non-laying.
7 months old when weighed 56 lb. Peaked at 75.4 lb
when 9 months old. Note very low mortality.
See notes of losses - autopsy, Cannibalism.
Fed TM-10 10/28/60 - 2 lbs?

Complete:

Comparison egg production house 4/59 (4th)
Moved house laid in 17 months 305 eggs each
beginning 10/59.

Dust count 12/60, 1/61, 2/61 - moult period.
3/61 - Thinned 11/61 - allow 6%
Culling = 282 hens
15% less production 2 yrs ago 57,501 45,684 13,185
9 mos prod. should have been

Allow 6% certain 1st year, 282 hens
got 57,501 - just yr - should have gotten 94,520

20,282 2nd yr 45,684
78,288 - less 9 eggs on floor 140,184
2 yrs 78,288
61,896

Note - no claim for losses in
overhead, hen damage, dairy, misc
feeds, etc